AD-A104 508

ROME AIR DEVELOPMENT CENTER GRIFFISS AFB NY
EFFECT OF ENERGETIC PARTICLE EVENTS ON VLF/LF PROPAGATION PARAM—ETC(U)
OCT 80 J P TURTLE, J E RASHUSSEN
RADC-TR-80-307

NL

Local Service Ser

AD A 104508

LEVEL (3) BS

RADC-TR-80-307 In-House Report October 1980

EFFECTS OF ENERGETIC PARTICLE EVENTS ON VLF/LF PROPAGATION PARAMETERS, 1974-1977

John P. Turtle John E. Rasmussen Wayne I. Klemetti



APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

L RIE COPY

ROME AIR DEVELOPMENT CENTER
Air Force Systems Command
Griffiss Air Force Base, New York 13441

This report has been reviewed by the RADC Public Affairs Office (PA) is releasable to the National Technical Information Service (MTIS). At 17 it will be releasable to the general public, including foreign nations.

RADC-TR-80-307 has been reviewed and is approved for publication.

APPROVED: Edward a. Lewis

EDWARD A. LEWIS, Chief Propagation Branch

Electromagnetic Sciences Division

APPROVED: Ofland School ALLAN C. SCHELL, Chief

Electromagnetic Sciences Division

FOR THE COMMANDER: Jako P. Kluss

JOHN P. HUSS

Acting Chief, Plans Office

If your address has changed or if you wish to be removed from the RADC meiling list, or if the addresses is no longer employed by your organization, please notify RADC (EEP) Hanscom AFB MA 01731. This will assist us in maintaining a current mailing list.

Do not return this copy. Retain or destroy.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

REPORT DOCUMENTATION PAGE		BEFORE COMPLETING FORM				
REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER				
RADC-TR-80-307	AD-1404508					
A TITLE (and Subusto)		5. TYPE OF REPORT & PERIOD COVERED				
EFFECTS OF ENERGETIC PA	ARTICLE EVENTS	In-House				
ON VLF/LF PROPAGATION	PARAMETERS,	in-nouse				
1974-1977.	-	6. PERFORMING ORG. REPORT NUMBER				
Z. AUTHOR(A)		8. CONTRACT OR GRANT NUMBER(4)				
John P. /Turtle						
John E./Rasmussen / Wayne I./Klemetti♥						
9. PERFORMING ORGANIZATION NAME AND ADD	Dec.	10. PROGRAM ELEMENT PROJECT TASK				
Deputy for Electronic Technol	1	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS				
Hanscom AFB	logy (RADC/EELE)	62702F				
Massachusetts 01731	10	4600 604				
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE				
Deputy for Electronic Technol		October 1980				
Hanscom AFB		19. NUMBER OF PAGES				
Massachusetts 01731	· 11/21	68 1-				
14. MONITORING AGENCY NAME & ADDRESS(II d.	ifferent from Controlling Office)	15. SECURITY CLASS, (of this report)				
		Unclassified / 61/				
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE				
16. DISTRIBUTION STATEMENT (of this Report)						
Approved for public release; of	distribution unlimite	d.				
•						
		1				
17. DISTRIBUTION STATEMENT (of the abstract or	the state of the state of the	- B				
17. DISTRIBUTION STATEMENT (OF the abstract en	ntered in Block 20, it dillerent from	(Report)				
18. SUPPLEMENTARY NOTES						
*Megapulse Inc., Bedford, M.	Α					
19. KEY WORDS (Continue on reverse side if necess	ary and identify by block number)					
VLF propagation						
LF propagation						
Ionospheric disturbances						
Polar cap absorption events						
20. ASSURACT (Continue on reverse side if necessa						
		re effects of energetic par-				
This report provides a summary of disturbance effects of energetic particle events on VLF/LF propagation parameters as observed by the USAF						
High Resolution VLF/LF Ionosounder in Northern Greenland. Disturbance						
effects on ionospheric reflectivity parameters, including reflection heights						
and coefficients, are presented along with data from a riometer, a magnetom-						
eter, and satellite particle detectors.						
√						
	<i>[[</i>]\`\`	ł				

DD 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20716

Preface

The authors thank Royce C. Kahler and Duane Marshall for help with the instrumentation which made the measurements possible, and Jens Ostergaard and Bjarne Ebbesen for the outstanding operation in Qanaq, Greenland.

Appreciation is also extended to the Danish Commission for Scientific Research in Greenland for allowing these measurements to be conducted, and to Jorgen Taagholt and V. Neble Jensen of the Danish Meteorological Institute's Ionospheric Laboratory for their continued cooperation in this program.

Accession For
NTIS COAST
DTIC TAX
Unennount of
Justification
Fy
Distribution
Avaitable and are
Special Contract
_
lack
R

1. INTRODUCTION 7 2. EVENT DATA 2.1 Observed Waveforms 13 2.2 Quantitative Reflection Parameters 14 2.2.1 Reflection Heights 14 2.2.2 Reflection Coefficients 14 2.3 Polarization Ellipses for the Down-Coming Skywaves 15 3. SUPPLEMENTARY DATA 15 4. DISTURBANCE CHARACTERISTICS 17 REFERENCES 67 Illustrations

1. Ionosounder Propagation Path, Thule AB - Qanaq, Greenland

2. Transmitting Antenna, Thule AB - Qanaq, Greenland

4. Orthogonal Receiving Antennas, Qanaq, Greenland

5. Example of Parallel and Perpendicular Waveforms

3. Basic Ionosounding Experiment

10

10

11

12

Contents

6.	. Fourier Amplitude Spectrum of Transmitted Pulse		
7.	Conversion Curve, Groundwave-Skywave Arrival Time Difference to Reflection Height	13	
8.	VLF/LF Ionospheric Reflectivity Data for 5 November 1974 (DOY 309) Solar Particle Event	19	
9.	VLF/LF Ionospheric Reflectivity Data for 30 April 1976 (DOY 121) Solar Particle Event	27	
10.	VLF/LF Ionospheric Reflectivity Data for 22 August 1976 (DOY 235) Solar Particle Event	35	
11.	VLF/LF Ionospheric Reflectivity Data for 26 July 1977 (DOY 207) Solar Particle Event	43	
12.	VLF/LF Ionospheric Reflectivity Data for 24 September 1977 (DOY 267) Solar Particle Event	51	
13.	VLF/LF Ionospheric Reflectivity Data for 22 November 1977 (DOY 326) Solar Particle Event	59	
		Tables	
1.	Disturbance Event Data	16	

Effects of Energetic Particle Events on VLF/LF Propagation Parameters, 1974—1977

1. INTRODUCTION

A compilation of data on the VLF/LF reflectivity of the polar ionosphere from 1975 to 1977 has been published in previous technical reports. $^{1-10}$ In this report, the data for specific periods are expanded in order to give a more detailed presentation of the effects of energetic particle events on VLF/LF propagation parameters. These periods have been chosen to show disturbance effects for events in which the 13.7 to 25.2 - MeV proton flux recorded by the IMP 7/8 satellites exceeded 10^{-2} particles/cm² sec sr MeV. The propagation data were obtained by the USAF High Resolution VLF/LF Ionosounder 11 , 12 which provides direct measurements of ionospheric reflection height and the reflection coefficient matrix elements $_{||}R_{||}$ and $_{||}R_{1}$. 13 Also included are data on particle flux density, HF riometer absorption, and geomagnetic field intensity.

(Received for publication 6 October 1980.)

Due to the number of references cited on this page, the reader is directed to pages 67 and 68 for references 1 through 10.

- 11. Lewis, E. A., Rasmussen, J. E., and Kossey, P. A. (1973) Measurements of ionospheric reflectivity from 6 to 35 kHz, J. Geophys. Res. 78:19.
- Kossey, P. A., Rasmussen, J. E., and Lewis, E. A. (1974) VLF pulse ionosounder measurements of the reflection properties of the lower ionosphere, <u>Akademie Verlag</u>, COSPAR.
- 13. Budden, K. G. (1961) Radio Waves in the Ionosphere, Cambridge University Press, London, p. 85.

The VLF/LF Ionosounding Transmitter (Figure 1) is located at Thule Air Base, Greenland (76° 33' N Lat., 68° 40' W Long.), and the receiving site is 106 km north at the Danish Meteorological Institute's Ionospheric Observatory in Qanaq, Greenland (77° 24' N Lat., 69° 20' W Long., Geomagnetic Lat. 89° 06' N). The ionosounding transmissions consist of a series of extremely short (approximately 100 usec) VLF pulses, precisely controlled in time, and radiated from the 130-m vertical antenna (Figure 2). At the receiver, the radiated signal arrives first by groundwave propagation (Figure 3). Due to the extremely short pulse length, this signal has passed the receiver before the arrival of the ionospherically reflected skywave pulse, providing independent groundwave and skywave data. Orthogonal loop antennas (Figure 4) are used to receive the two polarization components of the ionospherically reflected skywave signal. One loop, oriented in the plane of propagation, senses the groundwave and the unconverted or "parallel" (II) component of the down-coming skywave; the second loop, nulled on the groundwave. senses the converted or perpendicular (1) skywave component. The signal from each of the antennas is digitally averaged to improve the signal-to-noise ratio of the individual received waveforms before they are recorded on magnetic tape. An example of the observed waveforms is given in Figure 5, where the parallel waveform (a) consists of a groundwave propagated pulse, a quiet interval containing low level, off path groundwave reflections, followed by the first-hop parallel skywave component; the perpendicular waveform (b) is also shown. Each of these waveforms is comprised of 256 digitally averaged points, spaced 2 usec apart. Ionospheric reflection parameters are derived by computer processing of the ground and ionospherically reflected waveforms, with allowance made for factors such as ground conductivity and antenna patterns (see Section 2.2).

On 25 September 1975, and on 1 May 1977, the transmitted waveform was changed slightly; Figure 6 shows the resulting effect on the Fourier amplitude spectrum of the received groundwave signal. Although the data presented are generally limited to frequencies in the first, or principal lobe of the spectrum, information at higher frequencies can be used when sufficient signal-to-noise conditions exist. There is, however, a frequency range around each spectral null where insufficient signal exists for measurements.

2. EVENT DATA

The data are presented for each disturbance event in three general formats: first, the observed waveforms are shown in a synthetic three-dimensional display which starts approximately two days prior to the event and covers a fourteen-day period; second, the data are presented in the frequency domain with reflection




Figure 1. Ionosounder Propagation Path, Thule AB - Qanaq, Greenland

Figure 2. Transmitting Antenna, Thule AB - Qanaq, Greenland




Figure 3. Basic Ionosounding Experiment




Figure 4. Orthogonal Receiving Antennas, Qanaq, Greenland





Figure 5. Example of Parallel and Perpendicular Waveforms




heights and coefficients plotted as a function of frequency over the range from approximately 5 to 30 kHz; third, the data are presented as a function of time-of-day. In addition to reflection information, this section contains data on ionospheric absorption, geomagnetic field data activity, and solar proton fluxes.

2.1 Observed Waveforms

A three-dimensional waveform display is presented for a 2-week period containing each disturbance event, together with a display of the same 2-week period from a year in which it was not disturbed. For each display, the waveforms were stacked one behind the other in linear time, progressing from bottom to top. Each individual waveform is a 30-min average of approximately 10,000 pulses. The horizontal scale for these plots is linear in time (microseconds), measured from the start of the groundwave. This scale can be used to calculate an effective height of reflection by attributing the time delay between the start of the groundwave and the start of the skywave to a difference in travel distance, assuming a sharply bounded, mirror-like ionosphere. Figure 7 gives a conversion curve for this calculation based on simple geometry and the specific Thule AB - Qanaq, Greenland separation of 106 km. For the disturbance periods, fixed local ground clutter, amounting to only 2 percent of the groundwave amplitude, was removed to avoid interference with the skywave and improve the appearance of the waveforms.

Figure 7. Conversion Curve, Groundwave-Skywave Arrival Time Difference to Reflection Height



The three-dimensional displays of the disturbed and normal parallel waveforms are given for each event in Parts A and B of Figures 8 through 13. A plot
of the diurnal variation in solar zenith angle for the midpoint of the path appears
in Part C. The perpendicular waveform displays are shown in Parts D and E.
The time of maximum particle flux is indicated on the disturbance plots.

2.2 Quantitative Parallel and Perpendicular Reflection Parameters

For each event individual parallel and perpendicular waveforms were selected in order to show the effects of the disturbance on the ionospheric reflection height and reflection coefficients as a function of frequency. The selected waveforms from the disturbance period are shown in Part F of the data figures, whereas the corresponding undisturbed waveforms are shown in Part G.

2.2.1 REFLECTION HEIGHTS

The group mirror height (GMH) of reflection was obtained by determining the group delay of the skywave relative to the groundwave and attributing this difference to a difference in the propagation distance. The group delay can be defined as the rate of change of phase with frequency as discussed in Lewis et al. 11 For the GMH data presented in this report, a finite frequency difference of 1.0 kHz was used, and the corresponding phase difference as a function of frequency for the groundwave and both skywave signals was obtained by Fourier analysis of the respective pulses. The GMH calculations took into account ground conductivity (10⁻³ mho/m is assumed), with the Wait and Howe¹⁴ corrections applied. Group mirror heights for the parallel and perpendicular waveforms are plotted as a function of frequency in Parts H and I of Figures 8 through 13 for both normal and disturbed conditions. The GMHs are also presented as a function of time-of-day for the average frequency of 16.5 kHz. In Figures 8 through 13, Parts L and O, parallel and perpendicular reflection height information is given based on two-hour averaged data for the two-week period; Parts V and W show the 24-hour period of the event onset in greater detail, based on 5-min averaged data. These parts include a normal reflection height curve for reference purposes. Each point of the reference height curve is an average, by two-hour time blocks, for the 14-day normal period indicated.

2.2.2 REFLECTION COEFFICIENTS

Assuming that the ionosphere acts as a "mirror" at the GMH, we obtained plane wave reflection coefficients ¹³ by comparing the ratio of the skywave Fourier amplitude at a specific frequency to that of the groundwave, taking into account the

^{14.} Wait, J. R. and Howe, H. H. (1956) Amplitude and Phase Curves for Groundwave Propagation in the Band 200 Cycles per Second to 500 Kilocycles, Natl. Bureau of Standards, U. S. Circ. No. 574.

wave spreading, earth curvature, ground conductivity, path lengths, and antenna patterns including ground image effects.

The reflection coefficient $_{||}R_{||}$, obtained from analysis of the parallel skywave component, is plotted as a function of frequency for both normal and disturbed conditions in Part H. From the corresponding perpendicular skywave pulses, the coefficient $_{||}R_{||}$ was obtained; it appears as a function of frequency in Part I. The $_{||}R_{||}$ coefficient for 16 kHz is plotted as a function of time-of-day in Part M along with the averaged normal coefficient. As with the reflection heights, a more detailed $_{||}R_{||}$ coefficient plot, based on 5-min averaged data is shown in Part V. To show the variation in reflectivity as a function of frequency during the event, the reflection coefficients were calculated at 8 kHz, 16 kHz, and 22 kHz and are plotted in Part N as a function of time for the 14-day period. The corresponding reflection coefficient plots for $_{||}R_{||}$ are given in Parts P, Q, and W.

For certain coefficient data points, plotted as asterisks, the reflection coefficient appears without a corresponding GMH. For these particular data, only the skywave-groundwave ratios could be obtained since the skywaves were too weak to provide reliable group delay information. The reflection coefficients were estimated using a nominal GMH of 80 km in the calculations. These estimated coefficient values are included in the averages presented in Parts M, N, P, and Q, but the assumed heights are not used in the GMH averages.

2.3 Polarization Ellipses for the Down-Coming Skywaves

As described by Rasmussen et al, ¹⁵ the polarization ellipse of the skywave can be determined from the amplitudes of the parallel and perpendicular components and their phase difference. Each ellipse represents the locus of the tip of the rotation field vector as seen when looking in the direction of propagation of the down-coming skywave. The ellipses are drawn to a scale in which the incident wave amplitude is unity, and each division on the axis is 0.1. The direction of rotation is indicated by an arrow. Parts J and K of Figures 8 through 13 present polarization ellipse data as a function of frequency at 5 kHz intervals based on the selected disturbed and normal waveforms of Parts F and G, respectively.

3. SUPPLEMENTARY DATA

In order to interpret the effects of ionospheric disturbances on the VLF'. F ionosounding data, information from several geophysical sensors is presented. Parts R and S of Figures 8 through 13 present data from a magnetometer and a

^{15.} Rasmussen, J. E., et al., (1975) Low Frequency Wave-Reflection Properties of the Equatorial Ionosphere, AFCRL-TR-75-0615.

30-MHz riometer operated by RADC at Thule Air Base. The riometer, the conventional monitor of ionospheric disturbances, measures the signal level of cosmic radio noise passing through the ionosphere. The cyclical diurnal variation seen in the riometer signal level during undisturbed periods is caused primarily by the earth's daily rotation with respect to the extraterrestrial noise sources. A decrease in the received noise level represents an increase in the absorption resulting from enhanced ionization due to energetic particles. The absorption effects of energetic particle events are seen as an abrupt decrease in the riometer signal level followed by a gradual recovery to normal over a period of several days. The magnetometer data plotted are the horizontal (H) component of the polar magnetic field determined by a 3-axis fluxgate magnetometer at Thule Air Base. The magnetometer responds to the effects of polar ionospheric current systems related to disturbance events.

In addition to the information from the ground-based monitors, particle flux data are presented from the Applied Physics Laboratory of Johns Hopkins University experiments aboard the IMP 7 and 8 satellites.* These satellites are in roughly circular orbits at about 35 earth radii. The data presented in Parts T and U are hourly averages of differential flux levels for protons in two energy ranges: 0.97 to 1.85 MeV and 13.7 to 25.2 MeV. These particle data are most important for relating the VLF/LF ionosounder effects to the size of a particular disturbance.

The supplementary data are summarized in Table 1.

Table 1. Disturbance Event Data

	Date	Maximum 13.7-25,2 MeV Protons/cm ² sr sec MeV	Minimum 16 kHz ll Reflection Height (km)	30 MHz Riometer (dB) Absorption	Illumination Conditions
5	Nov 74 Day 309	1.3	63	<0. 5	nighttime
30	Apr 76 Day 121	6	58	3	daytime
22	Aug 76 Day 235	0.6	60	1.7	daytime
26	July 77 Day 207	0.02	70	<0.5	daytime
24	Sep 77 Day 267	2	57	2	daytime
22	Nov 77 Day 326	14	64	0.75	nighttime

^{*}Particle data obtained from the National Space Science Data Center, Greenbelt, MD

4. DISTURBANCE CHARACTERISTICS

From the events included in this report, the following VLF/LF disturbance patterns can be seen in the ionosounding data. The reflection heights for both the parallel and perpendicular components drop coincident with the influx of energetic particles. The level to which the reflection heights drop depends first, upon the magnitude of the particle flux, and second, upon the presence or absence of solar illumination during the event. The lowest reflection heights for a given particle flux are attained during a daytime event. As seen in Table 1, the 22 Aug 76 (DOY 235) polar daytime event with a 13 to 25 MeV proton flux of 0.6 particles resulted in a 16-kHz reflection height of 60 km; whereas the more energetic 22 Nov 77 (DOY 326) polar nighttime event with flux of 14 particles caused a drop in reflection height to only 64 km. During polar daytime events with continuous $\,$ D-region illumination, there is little if any diurnal height variation even if there are variations before and after the event. During polar nighttime events, a combination of particle ionization and weak noontime solar illumination can produce a diurnal variation during the event when none was present before or even after. For day-night events with the sun rising and setting, the disturbed reflection heights show a strong diurnal pattern, sometimes with larger variations than either before or after the event.

Reflection coefficients during energetic particle disturbances also behave differently, depending upon the solar illumination conditions. During a normal polar daytime period, with the sun continually above the horizon, reflection coefficients are quite variable and much lower than during a normal nighttime period. During a daytime particle event, reflection coefficients can actually increase with respect to normal conditions, particularly for 8 and 16 kHz. The coefficients show less diurnal variation during the event since the effects of particle ionization appear to override the effects of varying solar zenith angle. For the two strong daytime events, 30 Apr 76 (DOY 122) and 22 Aug 76 (DOY 235), several days after event maximum, the ionospherically reflected pulse became very weak for a period of approximately one day, resulting in reduced reflection coefficients before returning to normal conditions. This behavior is not associated with additional particle precipitation; it may be a part of the daytime recovery process for the particular geometry of the Thule AB - Qanaq path. Reflection coefficient behavior for nighttime and day-night disturbances is less complex, and similar in pattern to the reflection heights. During a polar nighttime event, the reflection coefficients decrease coincident with the particle influx and then recover steadily to normal. During day-night disturbed conditions, the reflection coefficients show an enhanced diurnal variation compared with variations prior to and following the event.

5 November 1974 Solar Particle Event

Date: 5 November 1974 DOY: 309

Report Figure: 8

Related Solar Flare: 5 November 1530 UT Start of Ionospheric Effects: 1600 UT

Time of Maximum 13-25 MeV Proton Flux: 2200 UT Maximum Flux: 1.3 particles/cm² sec sr MeV

Length of Particle Event: 3 days

Lowest Reflection Height (16 kHz): 63 km

Time of Lowest Height: 1730 UT 30 MHz Riometer Absorption: 0.5 dB Solar Zenith Angle Range: 93°-121° Illumination Conditions: nighttime

Typical of polar nighttime conditions, before the event the $_{||}R_{||}$ 16 kHz ionospheric reflection coefficient was large (0.5) and the reflection height high (85 km). The disturbance effects on the propagation parameters were quite clear: a rapid decrease followed by a gradual return to normal. As seen in Parts L-Q, before the event there was little diurnal variation in the parameters due to insufficient noontime ionizing radiation. During the event, however, an enhanced variation is noted; this was probably due to a combination of particle ionization, and a small amount of noontime solar radiation.

It is probable that a small particle event occurred between 12 and 14 November (DOY 316-318). This can be seen in the reflection height and coefficient data in Parts L-Q, as well as in the particle and magnetometer data in Parts R and U.




Figure 8. VLF/LF Ionospheric Reflectivity Data for 5 November 1974 (DOY 309) Schar Particle Event (continued)




Figure 8. VLF/LF Ionospheric Reflectivity Data for 5 November 1974 (DOY 309) Solar Particle Event (continued)






Figure 8. VLF/LF Ionospheric Reflectivity Data for 5 November 1974 (DOY 309) Solar Particle Event (continued)




J. SKYWAVE POLARIZATION ELLIPSES - DISTURBED CONDITIONS




K. SKYWAVE POLARIZATION ELLIPSES - NORMAL CONDITIONS

Figure 8. VLF/LF Ionospheric Reflectivity Data for 5 November 1974 (DON 309) Solar Particle Event (continued)



).).

Figure 8. VLF/LF Ionospheric Reflectivity Data for 5 November 1974 (DOY 309) Solar Particle Event (continued)






Figure 8. VLF/LF lonospheric Reflectivity Data for 5 November 1974 (DOY 309) Solar Particle Event (continued)

30 April 1976 Solar Particle Event

Date: 30 April 1976 DOY: 121

Report Figure: 9

Related Solar Flare: 30 April 2014 UT x-ray class: X2

Start of Ionospheric Effects: 2050 UT

Time of Maximum 13-25 MeV Proton Flux: 1 May 0300 UT

Maximum Flux: 6 particles/cm² sec sr MeV

Length of Particle Event: 5 days

Lowest 16 kHz Reflection Height: 58 km

Time of Lowest Reflection Height: 1 May 0015 UT

30 MHz Riometer Absorption: 3.5 dB Solar Zenith Angle Range: 63°-91° Illumination Conditions: daytime

As is typical of normal daytime conditions the $_{||}$ R $_{||}$ 16 kHz reflection coefficients (Part H) were much lower (0.08) than during normal polar nighttime conditions (0.5). The effects of the disturbance on the reflection coefficients were less clearly defined than during a nighttime event. As seen in Part N, the parallel reflection coefficient showed an increase during the first part of the event. This was followed by a relatively stable period with little diurnal variation for the next two days. On 3 May (DOY 124), two days after event particle maximum, the reflected skywave pulse became very weak for about a day (Parts A and D). This is seen in the reflection coefficients in Parts N and Q which were lower on 3 May than during event maximum, partic 'arly for 16 and 22 kHz. This was not related to an increase in particle flux, but may be part of the recovery process as associated with the particular geometry of the Thule Air Base - Qanaq path. Subsequent to 3 May, there was a gradual recovery over the next four days. The reflection heights during the event followed the usual pattern of abrupt drop followed by a gradual recovery; there was no diurnal height variation during this daytime event. The reflection heights returned to normal on 3 May (DOY 124), several days before the reflection coefficients.




Figure 9. VLF/LF Ionospheric Reflectivity Data for 30 April 1976 (DOY 121) Solar Particle Event (continued)




Figure 9. VLF/LF Ionospheric Reflectivity Data for 30 April 1976 (DOV 121) Solar Particle Event (continued)






Figure 9. VLF/LF Ionospheric Reflectivity Data for 30 April 1976 (DOY 121) Solar Particle Event (continued)



J. SKYWAVE POLARIZATION ELLIPSES - DISTURBED CONDITIONS



K. SKYWAVE POLARIZATION ELLIPSES - NORMAL CONDITIONS

25KHZ

Figure 9. VLF/LF Ionospheric Reflectivity Data for 30 April 1976 (DOY 121) Solar Particle Event (continued)




Figure 9. VLF/LF Ionospheric Reflectivity Data for 30 April 1976 (DOY 121) Solar Particle Event (continued)




Figure 9. VLF/LF Ionospheric Reflectivity Data for 30 April 1976 (1808 121) Solar Particle Event (continued)

22 August 1976 Solar Particle Event

Date: 22 August 1976 DOY: 235

Report Figure: 10

Related Solar Flare: 22 August 1214 UT - Asray class: M3

Start of Ionospheric Effects: 1230 UT

Time of Maximum 13-25 MeV Proton Flux: 1300 14 Maximum Flux: 0.6 particles/cm² sr sec MeV

Length of Particle Event: 3 days

Lowest 16 kHz Reflection Height: 60 km Time of Lowest Reflection Height: 1600 UT

30 MHz Riometer Absorption: 1.7 dB Solar Zenith Angle Range: 65°-93° Illumination Conditions: daytime

As with other polar daytime disturbance events, the R reflection coefficients at maximum particle flux (Part II) showed an increase for some frequencies and a decrease for other frequencies. After event maximum, the reflection coefficients were relatively steady for the next several days as compared with the coefficients before and after the event (Parts N and Q). In the latter part of the event, when the ionosphere had nearly recovered, the reflected skywave pulse became very weak for a period of about a day (Parts A and D). This same effect was seen in the April 1976 event (Figure 9). As seen in Parts N and Q on 24 August (DOY 237), the reflection coefficients were below the level at event maximum, particularly for 16 and 22 kHz. This again appears to be part of the disturbance recovery process rather than the result of an enhancement in the particle event itself. The reflection heights showed the usual daytime event pattern of an abrupt drop followed by a gradual recovery with no diurnal variations.




Figure 10. VLF/LF Ionospheric Reflectivity Data for 22 August 1976 (DOY 235) Solar Particle Event (continued)




Figure 10. VLF/LF Ionospheric Reflectivity Data for 22 August 1976 (DOY 235) Solar Particle Event (continued)






Figure 10. VLF/LF Ionospheric Reflectivity Data for 22 August 1976 (DOY 235) Solar Particle Event (continued)



J. SKYWAVE POLARIZATION ELLIPSES - DISTURBED CONDITIONS



K. SKYWAVE POLARIZATION ELLIPSES - NORMAL CONDITIONS

Figure 10. VLF/LF Ionospheric Reflectivity Data for 22 August 1976 (DOY 235) Solar Particle Event (continued)




Figure 10. VLF/LF lonospheric Reflectivity Data for 22 August 1976 (DOY 235) Solar Particle Event (continued)




Figure 10. VLF/LF lonospheric Reflectivity Data for 22 August 1976 (DOY 235) Solar Particle Event (continued)

26 July 1977 Solar Particle Event

Date: 26 July 1977 DOY: 207

Report Figure: 11

Related Solar Flare: No data

Start of Ionospheric Effects: 1600 UT

Time of Maximum 13-25 MeV Proton Flux: 2000 UT Maximum Flux: 0.02 particles/cm² sec sr MeV

Length of Particle Event: 3 days

Lowest 16 kHz Reflection Height: 70 km

Time of Lowest Reflection Height: 0100 UT 27 July

30 MHz Riometer Absorption: 0.5 dB Solar Zenith Angle Range: 58°-83° Illumination Conditions: daytime

This was the smallest particle event to be included in this report. Unlike other events reported here, the particle flux rose rather slowly to a maximum. As is typical of polar daytime events with continuous D-region illumination, the reflection height curves show a drop in height, followed by a gradual return to normal with no diurnal variation. The reflection coefficients which prior to the event were quite low and irregular, increased, and showed less variation during the event. The ionospheric reflection coefficients returned to normal by 29 July.




Figure 11. VLF/LF Ionospheric Reflectivity Data for 26 July 1977 (DON 207) Solar Particle Event (continued)




Figure 11. VLF/LF Ionospheric Reflectivity Data for 26 July 1977 (DOY 207) Solar Particle Event (continued)






Figure 11. VLF/LF Ionospheric Reflectivity Data for 26 July 1977 (DOY 207) Solar Particle Event (continued)



J. SKYWAVE POLARIZATION ELLIPSES - DISTURBED CONDITIONS



K. SKYWAVE POLARIZATION ELLIPSES - NORMAL CONDITIONS

25KHZ

Figure 11. VLF/LF Ionospheric Reflectivity Data for 26 July 1977 (DOY 207) Solar Particle Event (continued)




Figure 11. VLF/LF Ionospheric Reflectivity Data for 26 July 1977 (DOY 207) Solar Particle Event (continued)




Figure 11. VLF/LF Ionospheric Reflectivity Data for 26 July 1977 (DOY 207) Solar Particle Event (continued)

24 September 1977 Solar Particle Event

Date: 24 September 1977 DOY: 267

Report Figure: 12

Related Solar Flare: No data

Start of Ionospheric Effects: 0615 UT

Time of Maximum 13-25 MeV Proton Flux: 1000 UT

Maximum Flux: 2 particles/cm² sec sr MeV

Length of Particle Event: 6 days

Lowest 16 kHz Reflection Height: 57 km

Time of Lowest Reflection Height: 1400 UT

30 MHz Riometer Absorption: 2 dB Solar Zenith Angle Range: 75°-103° Illumination Conditions: Day-night

This energetic particle event occurred during an already disturbed period; a series of events had occurred, beginning on 8 September (DOY 251). No data are presented for these events as ionosounding records are incomplete. This was a day-night event with the sun rising and setting. Unlike polar daytime or nighttime events, there was an enhanced diurnal variation of the reflection heights and coefficients during the event. Solar radiation and particle ionization resulted in lower noontime reflection heights; the absence of solar radiation at night allowed the recombination of electrons, and thus a partial recovery toward normal reflection height conditions.

Figure 12. VLF/LF Ionospheric Reflectivity Data for 24 September 1977 (DOL JOI)

· Event (continued)




Figure 12. VLF/LF Ionospheric Reflectivity Data for 24 September 1977 (DOY 267) Solar Particle Event (continued)






Figure 12. VLF/LF Ionospheric Reflectivity Data for 24 September 1977 (DOY 267) Solar Particle Event (continued)



SKYWAVE POLARIZATION ELLIPSES - DISTURBED CONDITIONS



K. SKYWAVE POLARIZATION ELLIPSES - NORMAL CONDITIONS

Figure 12. VLF/LF Ionospheric Reflectivity Data for 24 September 1977 (DOY 267) Solar Particle Event (continued)




Figure 12. VI.F/LF Ionospheric Reflectivity Data for 24 September 1977 (DOY 267) Solar Particle Event (continued)




Figure 12. VLF/LF Ionospheric Reflectivity Data for 24 September 1977 (DOY 267) Solar Particle Event (continued)

W. PERPENDICULAR 16.0 KHz RELECTIVITY DATA - 5 MIN. AVERAGE FOR DAY OF EVENT ONSET

4. PARALLEL 16.0 KHz REFLECTIVITY DATA -5 MIN. AVERAGE FOR DAY OF EVENT ONSET

22 November 1977 Solar Particle Event

Date: 22 November 1977 DOY: 326

Report Figure: 13

Related Solar Flare: 0945 UT x-ray class: X1

Start of Ionospheric Effects: 1030 UT

Time of Maximum 13-25 MeV Proton Flux: 1600 UT

Maximum Flux: 14 particles/cm² sec sr MeV

Length of Particle Event: 8 days Lowest 16 kHz Reflection Height: 64

Time of Lowest Reflection Height: 1700 UT 30 MHz Riometer Absorption: 0.75 dB Solar Zenith Angle Range: 96°-124° Illumination Conditions: nighttime

Based on the high energy proton flux, this was the strongest event during the period of this report. Because this was a polar nighttime event, however, the magnitude of the effects on the VLF/LF parameters was less than would have occurred had there been solar illumination. The daytime 20 August 1976 event with only 0.6 particles/cm² sec sr MeV produced a lower reflection height of about 60 km. As is typical of a nighttime event, the disturbance effects on the reflection heights and coefficients (Parts L-Q) are well defined: an abrupt drop followed by a gradual recovery basically following the particle flux curve.




Figure 13. VLF/LF Ionospheric Reflectivity Data for 22 November 1977 (DOY 326) Solar Particle Event (continued)




Figure 13. VI.F/LF Ionospheric Reflectivity Data for 22 November 1977 (DOY 326) Solar Particle Event (continued)






Figure 13. VLF/LF Ionospheric Reflectivity Data for 22 November 1977 (DOY 326) Solar Particle Event (continued)

GROUP MIRROR REFLECTION HEIGHTS AND ASSOCIATED 118_REFLECTION COFFFICIENTS FROM PERPENDICULAR SKYWANE DATA



J. SKYWAVE POLARIZATION ELLIPSES - DISTURBED CONDITIONS



K. SKYWAVE POLARIZATION ELLIPSES - NORMAL CONDITIONS

Figure 13. VLF/LF Ionospheric Reflectivity Data for 22 November 1977 (DOY 326) Solar Particle Event (continued)

The second of the second of the complete of the second of




Figure 13. VLF/LF Ionospheric Reflectivity Data for 22 November 1977 (DOY 326) Solar Particle Event (continued)




Figure 13. VLF/LF Ionospheric Reflectivity Data for 22 November 1977 (DOY 326) Solar Particle Event (continued)

References

- Rasmussen, J. E., McLain, R. J., and Turtle, J. P. (1976) <u>VLF/LF</u> <u>Reflectivity of the Polar Ionosphere, 19 January - 2 March 1975</u>, AFCRL-<u>TR-76-0045</u>, AD A022674.
- 2. Rasmussen, J. E., McLain, R. J., and Turtle, J. P. (1976) VLF/LF
 Reflectivity of the Polar Ionosphere, 2 March 3 May 1975, RADC-TR-76146, AD A026465.
- Rasmussen, J. E., McLain, R. J., Turtle, J. P., and Klemetti, W. I. (1976) VLF/LF Reflectivity of the Polar Ionosphere, 4 May - 5 July 1975, RADC-TR-76-270, AD A034023.
- Rasmussen, J. E., McLain, R. J., Turtle, J. P., and Klemetti, W. I. (1976)
 <u>VLF/LF Reflectivity of the Polar Ionosphere</u>, 20 July 20 September 1975,
 <u>RADC-TR-76-327</u>, AD A036913.
- Rasmussen, J. E., McLain, R. J., Turtle, J. P., and Klemetti, W. I. (1976)
 <u>VLF/LF Reflectivity of the Polar Ionosphere</u>, 21 <u>September 1975 3 January 1976</u>, RADC-TR-76-378, AD A037794.
- 6. Rasmussen, J. E., Turtle, J. P., Pagliarulo, R. P., and Klemetti, W. I. (1977) VLF/LF Reflectivity of the Polar Ionosphere, 4 Janua. y 3 July 1976, RADC-TR-77-68, AD A040920.
- Rasmussen, J. E., Turtle, J. P., Pagliarulo, R. P., and Klemetti, W. I. (1977) VLF/LF Reflectivity of the Polar Ionosphere, 1 August 1976 – 1 January 1977, RADC-TR-77-141, AD A044050.
- 8. Rasmussen, J. E., Turtle, J. P., Pagliarulo, R. P., and Klemetti, W. I. (1977) VLF/LF Reflectivity of the Polar Ionosphere, 2 January - 30 April 1977, RADC-TR-77-251, AD A047238.
- 9. Rasmussen, J. E., Turtle, J. P., Pagliarulo, R. P., and Klemetti, W. I. (1977) VLF/LF Reflectivity of the Polar Ionosphere, 1 May 3 September 1977, RADC-TR-77-428, AD A053236.

- 10. Pagliarulo, R. P., Turtle, J. P., Rasmussen, J. E., and Klemetti, W. I. (1978) VLF/LF Reflectivity of the Polar Ionosphere, 4 September 31 December 1977, RADC-TR-78-95, AD A060918.
- 11. Lewis, E. A., Rasmussen, J. E., and Kossey, P. A. (1973) Measurements of ionospheric reflectivity from 6 to 35 kHz, <u>J. Geophys. Res.</u> 78:19.
- Kossey, P. A., Rasmussen, J. E., and Lewis, E. A. (1974) VLF pulse ionosounder measurements of the reflection properties of the lower ionosphere, <u>Akademie Verlag</u>, COSPAR.
- 13. Budden, K. G. (1961) Radio Waves in the Ionosphere, Cambridge University Press, London, p. 85.
- 14. Wait, J. R. and Howe, H. H. (1956) Amplitude and Phase Curves for Ground-wave Propagation in the Band 200 Cycles per Second to 500 Kilocycles, Natl. Bur. Stand. U. S. Circ. No. 574.
- 15. Rasmussen, J. E., et al., (1975) <u>Low Frequency Wave-Reflection Properties</u> of the Equatorial Ionosphere, AFCRL-TR-75-0615.